



Hortiphyte PK

Recommended Application Rates - FOLIAR SPRAY

OUTDOOR CROPS - Fruit & Hops

Apples - Apply at 3.0 L/Ha in 400-1000 L/Ha water. Apply at Petal Fall and repeat at 2-3 week intervals or as required.
Cane Fruit (Newly planted) - Apply at 4.0 L/Ha in 400-600 L/Ha. Apply when Primocanes 8-12" tall and at 3-4 week intervals or as required.
Cane Fruit (Established) - Apply at 4.0 L/Ha in 500-1000 L/Ha water. Apply at Green bud stage and at 3-4 week intervals or as required.
Grapes - Apply at 3.0 L/Ha in 250-400 L/Ha water. Apply at early fruit set and repeat at 3-4 week intervals or as required.
Hops - Apply at 2.5 L/Ha in 500-1000 L/Ha water. Apply once bines trained and repeat at 2 week intervals or as required.
Strawberries (Newly planted) - Apply at 4.0 L/Ha in 400-600 L/Ha water. Apply 2-3 weeks after planting and repeat at 3-4 week intervals or as required.
Strawberries (Established) - Apply at 4.0 L/Ha in 600-1000 L/Ha water. Apply at Green bud stage and repeat at 3-4 week intervals or as required.

OUTDOOR CROPS - Field Vegetables

Brassicas - Apply at 2.5 L/Ha in 200-400 L/Ha water. Apply at 3-4 leaf stage and repeat at 2-3 week intervals or as required.
Broad Beans - Apply at 2.0 L/Ha in 200-400 L/Ha water. Apply at early flowering and repeat at 2 week intervals or as required.
Carrots - Apply at 2.0 L/Ha in 200-400 L/Ha water. Apply 2-3 weeks after emergence and repeat at 2-3 week intervals or as required.
Onions & Leeks - Apply at 2.0 L/Ha in 200-400 L/Ha water. Apply when plants 8-10 cm tall and repeat at 2-3 week intervals or as required.
Peas - Apply at 2.5 L/Ha in 200-400 L/Ha water. Apply at early flowering and repeat at 2-3 week intervals or as required.
Potatoes - Apply at 2.5 L/Ha in 200-600 L/Ha water. Apply prior to rows meet and at 7-14 day intervals or as required.

OUTDOOR CROPS - Ornamentals

Ornamentals (Including Nursery Stock, Roses and Christmas Trees) - Apply 4.0 L/Ha in 600 - 1000 L/Ha water. Apply as soon as there is sufficient leaf canopy and repeat at 2-3 week intervals or as required.

PROTECTED CROPS

Strawberries - Apply 2 - 4 ml/L as a high volume foliar spray. Apply 2-3 weeks after planting or as soon as there is sufficient leaf canopy and repeat at 2-3 week intervals or as required.
Nursery Stock (including Pot Plants and Bedding Plants) - Apply 2-4 ml/L as a high volume foliar spray or as a root drench. Apply as soon as leaf canopy is sufficiently mature, repeat at 2-3 week intervals or as required.

Tank Mixing:

HortiPhyte PK is physically and chemically compatible with most pesticides. Check by mixing small amount in the correct ratio adding the product to the water followed by pesticide. Crops can be susceptible to scorch from mixtures of HortiPhyte and pesticides applied during periods of stress.

Caution:

Avoid spraying during high temperatures and bright sunlight. Keep out of reach of children. Avoid contact with skin, eyes or clothing. Wash exposed area with water.

Precautions:

Causes serious eye irritation.

Wear protective gloves / eye protection.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

If eye irritation persists: Get medical advice/attention.



WARNING